OFFICE OF NAVAL RESEARCH

Contract N00014-86-K-0043

TECHNICAL REPORT No. 86

Effect of Finite Size on Magnetoresistance

by

H. R. Lee, H. G. Oh and Thomas F. George

Prepared for Publication

in

Physical Review B

Departments of Chemistry and Physics State University of New York at Buffalo Buffalo, New York 14260

October 1988

Reproduction in whole or in part is permitted for any purpose of the United States Government.

This document has been approved for public release and sale; its distribution is unlimited.



REPORT DOCUMENTATION PAGE Form Approved OMB No. 0704-0188							
1a. REPORT SE	ECURITY CLASS	IFICATION		1b. RESTRICTIVE	MARKING\$		· L
Unclassified							
2a. SECURITY CLASSIFICATION AUTHORITY				3. DISTRIBUTION/AVAILABILITY OF REPORT			
A DECLASSIFICATION COMPLETA DING SCHEDUIT				Approved for public release; distribution			
2b. DECLASSIFICATION / DOWNGRADING SCHEDULE				\$			unlimited
4. PERFORMIN	G ORGANIZAT	ON REPORT NUMBE	R(S)	S. MONITORING ORGANIZATION REPORT NUMBER(S)			
4. PERFORMING ORGANIZATION REPORT NUMBER(S) UBUFFALO/DC/88/TR-86				1			
ĺ							
68. NAME OF PERFORMING ORGANIZATION 66. OFFICE SYMBOL				7a. NAME OF MONITORING ORGANIZATION			
Depts. Chemistry & Physics (If applicable)				ì			
State University of New York							
6c. ADDRESS	(City, State, and	d ZIP Code)			ty, State, and ZIP	Code)	
Fronczak Hall, Amherst Campus					Program		
Buffalo, New York 14260					incy Street		
					ı, Virginia		
8a. NAME OF ORGANIZA	FUNDING / SPO	INSORING	8b. OFFICE SYMBOL (If applicable)	9. PROCUREMENT	T INSTRUMENT ID	ENTIFICA	TION NUMBER
			(ii applicable)		Contract NOC	014-86	5-K-0043
	of Naval		<u></u>				
	City, State, and				UNDING NUMBER		
Chemistry Program 800 N. Quincy Street				PROGRAM ELEMENT NO.	PROJECT NO.	TASK NO	WORK UNIT ACCESSION NO.
Arlington, Virginia 22217					i	ł	
	lude Security C			<u> </u>	 	ــــــــــــــــــــــــــــــــــــــ	
(Size on Magnetor	resistance			
12. PERSONAL	L AUTHOR(S)						
•		H.R. Lee,	H.G. Oh and Tho	mas F. Georg	<u>şe</u>		
13a. TYPE OF	REPORT	13b. TIME C	OVERED TO	14. DATE OF REPO Octobe	PRT (Year, Month, er 1988	Day) 1	5. PAGE COUNT 17
16. SUPPLEME	NTARY NOTAT	ION Prepare	d for publication	n in Physics	al Review R		
1		rrepare	a for publication	on in injuice	II WEALEM D		
	COSATI	copre	I so cupier trover	Caral and a second		1 1 1 2 2 2 2 2	to Mark and to
17. FIELD	GROUP	SUB-GROUP	18. SUBJECT TERMS (MAGNETORES IS	STANCE:	DISORDER	a identin LED	by block number)
LIELD	GROUP	308-01007	FINITE SIZE	EFFECTS	COMPLICAT	י עידידעי	
			METALLIC SYS		QUANTUM	CORRÉC	ctions. (ind)
19 ARSTRACT	(Continue on	reverse if necessary	and identify by block n	umber)			
			re studied for m		tence in a d	licorde	ared metallic
system.	. Quantum	corrections	to the conductiv	vity are stro	ongly affect	ed by	the presence
of an i	in-plane m	agnetic field	in a thin film.	They are a	also affecte	d sign	nificantly by
9			quantum size. H			_	· · · ·
•			y due to both ef				
1			ld is found by t		_		
			sults can be obt				
							1.
merica]	l results	are presented	for the given i	inelastic sca	attering ler	igth.	Killing to the
}							
A 20 DISTRIBUT	TION / AVAIL AP	ILITY OF ABSTRACT		121 ARSTRACT CO	CURITY CLASSIFIC	ATION	
	SIFIED/UNLIMIT		RPT. DTIC USERS		Unclassifie		
	F RESPONSIBLE		2 2 032113	22b. TELEPHONE			OFFICE SYMBOL
	. David L			(202) 696-4			

Effect of Finite Size on Magnetoresistance

H. R. Lee, H. G. Oh and Thomas F. George Department of Physics and Astronomy
 239 Fronczak Hall
 State University of New York at Buffalo
 Buffalo, New York 14260

C. I. Um
Department of Physics
College of Science
Korea University
Seoul 136, KOREA

Abstract

Finite size effects are studied for magnetoresistance in a disordered metallic system. Quantum corrections to the conductivity are strongly affected by the presence of an in-plane magnetic field in a thin film. They are also affected significantly by the boundaries of the finite quantum size. Expressions are obtained for the quantum correction to the conductivity due to both effects. The dephasing characteristic time scale due to the magnetic field is found by the exact eigenvalues of the system.

One-, two- and three-dimensional results can be obtained with the proper limits. Some numerical results are presented for the given inelastic scattering length.

PACS numbers: 72.15.Gd, 73.50.Dn, 73.60.Aq



An ediction does a service ser

I. <u>Introduction</u>

The effect of a magnetic field on electronic states has been studied extensively for disordered systems. 1-3 The predictions of the anomalous magnetoresistance have been tested by several experiments on metal films4 and semiconductor structures. 5 The resistance of thin films and wires has been studied in the presence of a longitudinal magnetic field, 3 which is more effective than a field directed perpendicular to the film. In Ref. 3, the dimensions of the system are very small compared to the magnetic length because the magnetic field is treated as a small perturbation. Here perturbation theory yields the dephasing characteristic time $\tau_{\rm H}$ - $12a_{\rm H}^2/{\rm DW}^2$ for a square film or a wire of rectangular cross section, where $\mathbf{a}_{\mathbf{H}}$ is the magnetic length of a particle with charge 2e, D is the electon diffusion coefficient, and W is the dimension of the system. Generally perturbation theory can be used when the condition $W << a_u$ holds. But, it is not appropriate to apply perturbation theory to the ground state, which is most important for corrections to the conductivity. The reason is explained later in this paper. Here we study the effect of the boundaries on the eigenvalues of the maximally crossed diagram in a thin film or wire with the longitudinal magnetic field by a numerical method. From the eigenvalues of the maximally crossed diagram, we obtain analytical expressions for quantum corrections to the conductivity of thin films and wires, which are given in Eqs. (13) and (14), respectively. Quantum corrections to the conductivity are calculated as a function of the normalized dimension $(W/a_{\rm H})$ of the thin film or wire. Since a wide range of the normalized dimension W/aH is used in our calculations, the actual dimension W of the system can range from

very small values to values which are larger than the magnetic length. We obtain the dephasing characteristic time $\tau_{\rm H}$ = 24 ${\rm a_H^2/DW^2}$ as different from the result of the perturbation method in restricted geometries.

II. Theory

The quantum correction to the conductivity of non-interacting electrons weakly scattered by rigid random impurities is

$$\Delta\sigma(\omega\to 0) = -\frac{\mathrm{se}^2}{\pi \mathrm{ld}} \mathrm{D}\Gamma(\vec{r},\vec{r}',\omega) \quad , \tag{1}$$

where s is spin degeneracy, and $\Gamma(\vec{r},\vec{r}',\omega)$ is the vertex correction due to the sum of all the maximally crossed diagrams. In the absence of the magnetic field, the vertex part $\Gamma(\vec{r},\vec{r}',\omega)$ is

$$\Gamma(\vec{r},\vec{r}',\omega) = \frac{1}{\mu} \sum_{\vec{q}} \frac{1}{Dq^2 + i\omega} . \qquad (2)$$

This vertex part is strongly affected by the presence of an external magnetic field, because the symmetries inherent to the system are broken by the field. This has been studied in Ref. 1 in the coordinate representation through the equation

$$N[D(-i\frac{\partial}{\partial \vec{r}} - \frac{2e}{Nc}\vec{A}(\vec{r}))^2 + \frac{1}{r_{in}}] \Gamma(\vec{r}, \vec{r}') = \delta(\vec{r} - \vec{r}') , \qquad (3)$$

where $\vec{A}(\vec{r})$ is the magnetic vector potential, and $-i\omega$ is replaced by the inelastic scattering time r_{in}^{-1} at finite temperature.

Now let us consider a thin film which has a thickness W under an inplane magnetic field $\vec{H} = (0,0,H_0) = \vec{\nabla} \times \vec{A}$. If we choose the Landau gauge $\vec{A} = (0, xH_0, 0)$, the solution of Eq. (3) can be written as

$$\Gamma(x,x') = \int \frac{dq_y dq_z}{(2\pi)^2 N} \sum_{n} \frac{\psi_{n,q_y}(x)\psi_{n,q_y}(x')}{D(q_z^2 + E_n(q_y)) + \frac{1}{r_{in}}}.$$
 (4)

where $\psi_{n,q_X}(x)$ and $E_n(q_y)$ are the eigenfunctions and eigenvalues, respectively, of the equation

$$- \left[\frac{\partial^2}{\partial x^2} - (q_y - \frac{2e}{\mu c} H_0 x)^2 \right] \psi_{n,q_y}(x) - E_n(q_y) \psi_{n,q_y}(x)$$
 (5)

within the film. The above equation can be written as

$$-\left[\frac{\partial^{2}}{\partial x^{2}} - \frac{(x-x_{0})^{2}}{a_{H}^{4}}\right] \psi_{n,q_{y}}(x) - E_{n}(q_{y})\psi_{n,q_{y}}(x) , \qquad (6)$$

where $a_H = \sqrt{\frac{c M}{2e H_0}}$ is the magnetic length of a doubly-charged particle and x_0 is related to the wavevector q_y by the expression $x_0 = a_H^2 q_y$. Now if we introduce the normalized coordinate $\xi = \sqrt{2}x/a_H$, Eq. (6) can be transformed into the well-known Weber equation,

$$\left[\frac{\partial^{2}}{\partial \xi^{2}} - \frac{1}{4}(\xi - \xi_{0})^{2} + (\nu_{n} + \frac{1}{2})\right] \psi_{\nu_{n}, q_{y}}(\xi) = 0 , \qquad (7)$$

where ξ_0 is related to the wavevector q_y by $\xi_0 = \sqrt{2}a_H q_y$. We construct the general solutions of Eq. (7) as

$$\psi_{\nu_n, q_y}(\xi) = A D_{\nu_n}(\xi) + D D_{\nu_n}(-\xi)$$
, (8)

where A and B are the normalization constants and $D_{\underline{m}}(z)$ is the Weber function given by

$$D_{\mathbf{m}}(z) = 2^{\mathbf{m}/2} \exp(-\frac{z^{2}}{4}) \left[\frac{\sqrt{\pi}}{\Gamma(-\frac{\mathbf{m}}{2} + \frac{1}{2})} {}_{1}F_{1}(-\frac{\mathbf{m}}{2}; \frac{1}{2}; \frac{1}{2}z^{2}) \right] - \frac{\sqrt{2\pi} z}{\Gamma(-\frac{\mathbf{m}}{2})} {}_{1}F_{1}(-\frac{\mathbf{m}}{2} + \frac{1}{2}; \frac{3}{2}; \frac{1}{2}z^{2}) \right] .$$
 (9)

Here $_1F_1(a;b;x)$ is the confluent hypergeometric function 8 and $\Gamma(z)$ is the gamma function. The eigenvalues are given by

$$E_n(q_y) = \frac{2}{a_H^2} (\nu_n + \frac{1}{2})$$
 (10)

III. Results and Discussion

Equation (7) yields the discrete spectrum of eigenvalues for each value of the continuously varying wavevector $\mathbf{q}_{\mathbf{v}}$. If there are no boundaries, both

the eigenvalues and eigenvectors become identical to the solutions describing the unrestricted motion of free particles in the magnetic field. The eigenvalues of Eq. (7) can be determined if the precise form of the confining potential is given. The results are shown in Fig. 1 for normalized dimensions of the sample $\sqrt{2}W/a_{H}^{-}$ 1.0 with the boundaries of an infinite confining potential barrier, given by

$$\frac{\partial \psi_{n,q_y}(x)}{\partial x} \Big|_{x=\frac{W}{2}} = 0 .$$
 (11)

Since each mode in Fig. 1 shows parabolic-like behavior as a function of the wavevector $\mathbf{q}_{\mathbf{y}}$, we may write the eigenvalues in a parabolic approximation as

$$E_{n}(q_{y}) = \frac{\Delta_{n}}{a_{H}^{2}} + c_{n}q_{y}^{2} , \qquad (12)$$

where Δ_n is the y-intercept $(q_y=0)$ and C_n represents the coefficient of the quadratic term in each mode.

For a thin film, when we substitute Eq. (12) into (4) and integrate over $\mathbf{q}_{\mathbf{y}}$ and $\mathbf{q}_{\mathbf{z}}$, we obtain

$$\Delta \sigma = \frac{e^2}{2\pi^2 N} \sum_{n} \frac{1}{\sqrt{C_n}} \ln \left(\frac{1/\ell_{e1}^2 + 1/\ell_{in}^2}{\Delta_{n}/a_H^2 + 1/\ell_{in}^2} \right) , \qquad (13)$$

where the elastic diffusion length $\ell_{\rm el} = \sqrt{Dr}_{\rm el}$ is used for the upper limit of the integration. The quantum corrections to the conductivity of rectangular wires with transverse dimensions small in comparison with $\ell_{\rm in}$ can be obtained in the same way and are given by

$$\Delta \sigma = -\frac{2e^2}{\pi^2 H} \sum_{n} \frac{1}{\sqrt{c_n} (\frac{1}{\ell_{in}^2} + \frac{\Delta_n}{a_H^2})} \tan^{-1} \left(\frac{c_n (1/\ell_{e1}^2 - \Delta_n/a_H^2)}{1/\ell_{in}^2 + \Delta_n/a_H^2} \right)^{l_i} . \tag{14}$$

Numerical results for magnetoconductivity in thin films and the magnetoconductivity per unit length in rectangular wires are presented in Figs. 2 and 3, respectively, for various values of the ratio W/ℓ_{in} . In each graph we have used the unitless quantity $\xi = \sqrt{2}W/a_H$ in the x-direction, and the dimensions of the sample are normalized to the inelastic scattering length ℓ_{in} . Thus x-coordinates of the graphs are proportional to $\sqrt{H_0}$ for the given sample size. The effect of a magnetic field becomes more prominant on the quantum interference (weak localization) at small values of W/ℓ_{in} , that is, at small dimensions of the sample or at large values of ℓ_{in} (low temperature) for the given system.

The values Δ_0 and C_0 of the lowest-lying state (n=0), which is the most important for quantum corrections to the conductivity, are given in Table I for comparison with those in the absence of a magnetic field. Without the magnetic field, $E_n^0(q_y)$ may be written in a similar form as

$$E_{n}^{0}(q_{y}) = (\frac{n\pi}{V})^{2} + q_{y}^{2} , \qquad (15)$$

where n = 0, ± 1 , ± 2 , . . . When the condition W << a_H holds, Table I shows $\Delta_0/a_H^2 = W^2/24a_H^4$ for the ground state, whereas perturbation theory yields $W^2/12a_H^4$. As we can see clearly from Eq. (5) that perturbation theory can not be applied for small values of $q_y \approx 0$, which has the most significant contributions to the corrections to the conductivity. Thus the characteristic time scale must be $\tau_H = 24a_H^2/DW^2$ in magnetoconductivity, which is given by

$$\sigma(H) - \sigma(0) = \frac{e^2}{2\pi^2 H} \ln(\frac{r_{in}}{r_H} + 1)$$
 (16)

for a thin film, and the quantum corrections to the conductivity per unit length are given by

$$\Delta\sigma(H) = -\frac{e^2}{\pi H} \left(\frac{1}{Dr_{in}} + \frac{1}{Dr_{H}} \right) \tag{17}$$

for a wire of rectangular cross section.

If the other condition W >> a_H holds, the eigenvalues are divided into two parts: (1) the surface part, that is, those states whose orbit is affected by one wall of the sample, and (2) the bulk orbits which are not affected by the boundaries of the sample. The surface part always has an almost constant contribution to the conductivity, whereas the bulk part is proportional to the sample width W due to many degenerate states in each

level. Thus, we can recover the results of the bulk limit given in Ref. 1 (cf. the values in Table I).

This research was supported by the Office of Naval Research and the Air Force Office of Scientific Research (AFSC), United States Air Force, under Contract No. F49620-86-C-0009. The United States Government is authorized to reproduce and distribute eprints for governmental purposes notwithstanding any copyright notation hereon.

References

- B. L. Al'tshuler, D. Khemel'nitzkii, A. I. Larkin, and P. A. Lee, Phys. Rev. B 22, 5142(1980)
- 2. P. A. Lee and T. V. Ramakrishnan, Rev. Mod. Phys. <u>57</u>, 287 (1985).
- 3. B. L. Al'tshuler, and A. G. Aronov, JETP Lett. 33, 499 (1981).
- J. J. Lin, and N. Giordano, Phys. Rev. B <u>33</u>, 1519 (1986); S. Wind, M. J. Rooks, V. Chandrasekhar and D. E. Prober, Phys. Rev. Lett. <u>57</u>, 633 (1986).
- R. G. Wheeler, K. K. Choi, A. Goel, R. Wisnieff and D. E. Prober, Phys.
 Rev. Lett. 49, 1674 (1982); T. J. Thornton, M. Pepper, H. Ahmed, D.
 Andrews and G. J. Davies, ibid. 56, 1198 (1986)
- D. J. Thouless, Phys. Rev. Lett. <u>39</u>, 1167(1977); P. W. Anderson, E.
 Abrahams and T. V. Ramakrishnan, Phys. Rev. Lett. <u>43</u>, 718 (1979).
- 7. P. M. Morse, and H. Feshbach, <u>Methods of Theoretical Physics</u> (McGraw-Hill, New York, 1953), p. 1403.
- 8. M. Abramowitz, and I. A. Stegun, <u>Handbook of Mathematical Functions</u>.

 (Dover, New York, 1964), p. 503.

Rigenvalue shift (Δ_0) and parabolic coefficient (C_0) due to the boundaries and the longitudinal magnetic field. TABLE I.


£(W)	$\xi(W)$ 1.0×10 ⁻⁴ 1.0×10 ⁻³	1.0×10 ⁻³		0.1	1.0	2.0	3.0	4.0
٥	4.167×10 ⁻⁹	4.167×10 ⁻⁹ 4.167×10 ⁻⁷		4.167×10^{-5} 4.167×10^{-3} 4.165×10^{-2} 1.656×10^{-1} 3.631×10^{-1} 6.018×10^{-1}	4.165×10 ⁻²	1.656×10 ⁻¹	3.631×10 ⁻¹	6.018×10
,	1.000	1.000	1.000	1.000	0.992	0.912	0.797	0.687


TABLE I. Eigenvalue shift (Δ_0) and parabolic coefficient (C_0) due to the boundaries and the longitudinal magnetic field.


$$\xi$$
(W) 1.0 × 10⁻⁴ 1.0 × 10⁻³ 1.0 × 10⁻² 0.1 1
 Δ_0 4.167 × 10⁻⁹ 4.167 × 10⁻⁷ 4.167 × 10⁻⁵ 4.167 × 10⁻³ 4.165
 c_0 1.000 1.000 1.000 0.

Figure Captions

- 1. The energy dispersion in a longitudinal magnetic field with dimensions of the system $\frac{\sqrt{2}W}{a_H} = 1.0$ and infinite-barrier confining potential. The x-coordinate ξ_0 is related to the wavevector q_y by $\xi_0(q_y) = \sqrt{2}a_Hq_y$.
- 2. The magnetoconductivity (divided by the coefficient $\frac{e^2}{2\pi^2 k}$) of a thin film plotted against the unitless parameter $\xi(a_H) = \frac{\sqrt{2}W}{a_H}$. (1) $W/l_{in} = 0.05$, (2) $W/l_{in} = 0.2$ and (3) $W/l_{in} = 1.0$.
- 3. The magnetoconductivity (divided by the coefficient $\frac{2e^2 \ell_{in}}{\pi^2 k}$) per unit length of a rectangular wire plotted against the unitless parameter $\xi(a_H) = \frac{\sqrt{2}W}{a_H}$. (1) W/ $\ell_{in} = 0.05$, (2) W/ $\ell_{in} = 0.2$ and (3) W/ $\ell_{in} = 1.0$.







01/1113/86/2

TECHNICAL REPORT DISTRIBUTION LIST, GEN

:	No. Copies	•	No. Copies
Office of Naval Research Attn: Code 1113 800 N. Quincy Street Arlington, Virginia 22217-5000	2	Dr. David Young Code 334 NORDA NSTL, Mississippi 39529	1
Dr. Bernard Douda Naval Weapons Support Center Code 50C Crane, Indiana 47522-5050	1	Naval Weapons Center Attn: Dr. Ron Atkins Chemistry Division China Lake, California 93555	1
Naval Civil Engineering Laboratory Attn: Dr. R. W. Drisko, Code L52 Port Hueneme, California 93401	1	Scientific Advisor Commandant of the Marine Corps Code RD-1 Washington, D.C. 20380	1
Defense Technical Information Center Building 5, Cameron Station Alexandria, Virginia 22314	12 high quality	U.S. Army Research Office Attn: CRD-AA-IP P.O. Box 12211 Research Triangle Park, NC 27709	1
DTNSRDC Attn: Dr. H. Singerman Applied Chemistry Division Annapolis, Maryland 21401	1	Mr. John Boyle Materials Branch Naval Ship Engineering Center Philadelphia, Pennsylvania 19112	1
Dr. William Tolles Superintendent Chemistry Division, Code 6100 Naval Research Laboratory	1	Naval Ocean Systems Center Attn: Dr. S. Yamamoto Marine Sciences Division San Diego, California 91232	1
Washington, D.C. 20375-5000	i	Dr. David L. Nelson Chemistry Division Office of Naval Research 800 North Quincy Street Arlington, Virginia 22217	1

Dr. J. E. Jensen Hughes Research Laboratory 3011 Malibu Canyon Road Malibu, California 90265

Dr. J. H. Weaver
Department of Chemical Engineering
and Materials Science
University of Minnesota
Minneapolis, Minnesota 55455

Dr. A. Reisman Microelectronics Center of North Carolina Research Triangle Park, North Carolina 27709

Dr. M. Grunze
Laboratory for Surface Science and
Technology
University of Maine
Orono, Maine 04469

Dr. J. Butler Naval Research Laboratory Code 6115 Washington D.C. 20375-5000

Dr. L. Interante Chemistry Department Rensselaer Polytechnic Institute Troy, New York 12181

Dr. Irvin Heard Chemistry and Physics Department Lincoln University Lincoln University, Pennsylvania 19352

Or. K.J. Klaubunde Department of Chemistry Kansas State University Manhattan, Kansas 66506 Dr. C. B. Harris
Department of Chemistry
University of California
Berkeley, California 94720

Dr. F. Kutzler
Department of Chemistry
Box 5055
Tennessee Technological University
Cookesville, Tennessee 38501

Dr. D. Dilella Chemistry Department George Washington University Washington D.C. 20052

Dr. R. Reeves Chemistry Department Renssaeler Polytechnic Institute Troy, New York 12181

Dr. Steven M. George Stanford University Department of Chemistry Stanford, CA 94305

Dr. Mark Johnson Yale University Department of Chemistry New Haven, CT 06511-8118

Dr. W. Knauer Hughes Research Laboratory 3011 Malibu Canyon Road Malibu, California 90265

Cr. G. A. Somorjai Department of Chemistry University of California Berkeley, California 94720

Dr. J. Murday Naval Research Laboratory Code 6170 Washington, D.C. 20375-5000

Dr. J. B. Hudson Materials Division Rensselaer Polytechnic Institute Troy, New York 12181

Dr. Theodore E. Madey
Surface Chemistry Section
Department of Commerce
National Bureau of Standards
Washington, D.C. 20234

Dr. J. E. Demuth
IBM Corporation
Thomas J. Watson Research Center
P.O. Box 218
Yorktown Heights, New York 10598

Dr. M. G. Lagally
Department of Metallurgical
and Mining Engineering
University of Wisconsin
Madison, Wisconsin 53706

Dr. R. P. Van Duyne Chemistry Department Northwestern University Evanston, Illinois 60637

Dr. J. M. White Department of Chemistry University of Texas Austin, Texas 78712

Dr. D. E. Harrison Department of Physics Naval Postgraduate School Monterey, California 93940 Dr. R. L. Park
Director, Center of Materials
Research
University of Maryland
College Park, Maryland 20742

Dr. W. T. Peria Electrical Engineering Department University of Minnesota Minneapolis, Minnesota 55455

Dr. Keith H. Johnson
Department of Metallurgy and
Materials Science
Massachusetts Institute of Technology
Cambridge, Massachusetts 02139

Dr. S. Sibener Department of Chemistry James Franck Institute 5640 Ellis Avenue Chicago, Illinois 60637

Dr. Arnold Green Quantum Surface Dynamics Branch Code 3817 Naval Weapons Center China Lake, California 93555

Dr. A. Wold Department of Chemistry Brown University Providence, Rhode Island 02912

Dr. S. L. Bernasek Department of Chemistry Princeton University Princeton, New Jersey 08544

Dr. W. Kohn
Department of Physics
University of California, San Diego
La Jolla, California 92037

Dr. F. Carter Code 6170 Naval Research Laboratory Washington, D.C. 20375-5000

Dr. Richard Colton Code 6170 Naval Research Laboratory Washington, D.C. 20375-5000

Dr. Dan Pierce National Bureau of Standards Optical Physics Division Washington, D.C. 20234

Dr. R. Stanley Williams Department of Chemistry University of California Los Angeles, California 90024

Dr. R. P. Messmer Materials Characterization Lab. General Electric Company Schenectady, New York 22217

Dr. Robert Gomer Department of Chemistry James Franck Institute 5640 Ellis Avenue Chicago, Illinois 60637

Dr. Ronald Lee R301 Naval Surface Weapons Center White Oak Silver Spring, Maryland 20910

Dr. Paul Schoen Code 6190 Naval Research Laboratory Washington, D.C. 20375-5000 Dr. John T. Yates Department of Chemistry University of Pittsburgh Pittsburgh, Pennsylvania 15260

Dr. Richard Greene Code 5230 Naval Research Laboratory Washington, D.C. 20375-5000

Dr. L. Kesmodel
Department of Physics
Indiana University
Bloomington, Indiana 47403

Dr. K. C. Janda University of Pittsburg Chemistry Building Pittsburg, PA 15260

Dr. E. A. Irene
Department of Chemistry
University of North Carolina
Chapel Hill, North Carolina 27514

Dr. Adam Heller Bell Laboratories Murray Hill, New Jersey 07974

Dr. Martin Fleischmann Department of Chemistry University of Southampton Southampton 509 5NH UNITED KINGDOM

Dr. H. Tachikawa Chemistry Department Jackson State University Jackson, Mississippi 39217

Dr. John W. Wilkins Cornell University Laboratory of Atomic and Solid State Physics Ithaca. New York 14853

Dr. R. G. Wallis Department of Physics University of California Irvine, California 92664

Dr. D. Ramaker Chemistry Department George Washington University Washington, D.C. 20052

Dr. J. C. Hemminger Chemistry Department University of California Irvine, California 92717

Dr. T. F. George Chemistry Department University of Rochester Rochester, New York 14627

Dr. G. Rubloff IBM Thomas J. Watson Research Center P.O. Box 218 Yorktown Heights, New York 10598

Dr. Horia Metiu Chemistry Department University of California Santa Barbara, California 93106

Dr. W. Goddard
Department of Chemistry and Chemical
Engineering
California Institute of Technology
Pasadena, California 91125

Dr. P. Hansma
Department of Physics
University of California
Santa Barbara, California 93106

Dr. J. Baldeschwieler
Department of Chemistry and
Chemical Engineering
California Institute of Technology
Pasadena, California 91125

Dr. J. T. Keiser Department of Chemistry University of Richmond Richmond, Virginia 23173

Or. R. W. Plummer Department of Physics University of Pennsylvania Philadelphia, Pennsylvania 19104

Dr. E. Yeager Department of Chemistry Case Western Reserve University Cleveland, Ohio 41106

Dr. N. Winograd
Department of Chemistry
Pennsylvania State University
University Park, Pennsylvania 16802

Dr. Roald Hoffmann Department of Chemistry Cornell University Ithaca, New York 14853

Dr. A. Steckl
Department of Electrical and
Systems Engineering
Rensselaer Polytechnic Institute
Troy, NewYork 12181

Dr. G.H. Morrison Department of Chemistry Cornell University Ithaca, New York 14853